
TO FILE (1) AD-A214 674

IDENTIFYING THE CIRCADIAN
CYCLE IN HUMAN INFORMATION
PROCESSING DATA USING
PERIODICITY ANALYSIS:

A SYNOPSIS

Diane L. Damos

DTIC ELECTE NOV 3 0 1989



89 11 27 042

Naval Aerospace Medical Research Laboratory

Naval Air Station

Pensacola, Florida 32508-5700

Approved for public release; distribution unlimited.

Reviewed and approved 3 april 1989

A. BRADY, CAPT, ASC USN Commanding Officer



This research was sponsored by the Naval Medical Research and Development Command under work unit 62758N MF58528 MF5852801B 01B.

The views expressed in this article are those of the author and do not reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

Trade names of materials and/or products of commercial or nongovernment organizations are cited as needed for precision. These citations do not constitute official endorsement or approval of the use of such commercial materials and/or products.

Reproduction in whole or in part is permitted for any purpose of the United States Government.

		REPORT D	OCUMENTATIO	I PAGE			Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION				1b. RESTRICTIVE MARKINGS				
Unclassified				D. DISTRIBUTION ANALIABILITY OF DEPOST				
2a. SECURITY CLASSIFICATION AUTHORITY				3. DISTRIBUTION/AVAILABILITY OF REPORT				
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE				Approved for public release; distribution unlimited.				
4. PERFORMIN	G ORGANIZATIO	ON REPORT NUMBE	R(S)	5. MONITORING	ORGANIZATION RE	PORT NUMBER	R(S)	
NAMRL Tec	hnical Mem	orandum 89-1	•				,	
6a. NAME OF PERFORMING ORGANIZATION 6b. Naval Aerospace Medical			6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION				
Naval Aerospace Medical Research Laboratory			00	Naval Medical Research and Development Command			elobment	
				7b. ADDRESS (City, State, and ZIP Code)				
6c. ADDRESS (	Station	2,, 1001,		Naval Medical Command				
	, FL 32508	3-5700		National Capital Region Bethesda, MD 20814-5044				
8a. NAME OF	FUNDING / SPOR	NSORING	8b. OFFICE SYMBOL	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER				
ORGANIZA NMRDC	ATION		(If applicable) Code 404					
8c. ADDRESS (	City, State, and	ZIP Code) NMRDC, Nava		10. SOURCE OF FUNDING NUMBERS				
		, NMRDC, Nava Capital Regio		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.	
	MD 20814-		, , , , , , , , , , , , , , , , , , ,	62758N	MF58528	01B		
12. PERSONAI Damos, Di	LAUTHOR(S)	alysis: A Sy	-	TA DATE OF DEDO	DT /Vaar Manth /	131) 115 PA	SE COUNT	
13a. TYPE OF Interim	REPORT	13b. TIME C	TO	14. DATE OF REPORT (Year, Month, Day) 15. PAGE COUNT 89-04-03				
16. SUPPLEMI	ENTARY NOTAT	ION				<u>-</u> . –		
17.	COSATI	CODES	18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)					
FIELD	GROUP	\$UB-GROUP	Circadian ef	•	ormation pro			
			Statistical.	tests, pe	eformance (	humam)	(KT)	
Data of many o	from human	n information iodicity tech	and identify by block on processing ta hniques. This of the assumpti	sks frequent memorandum i ons of the m	dentifies a lore familia	ppropriat	e techniques	
	TION / AVAILABI	ILITY OF ABSTRACT	RPT	21. ABSTRACT SECURITY CLASSIFICATION Unclassified				
22a. NAME C	OF RESPONSIBLE	INDIVIDUAL	manding Officer	22b. TELEPHONE	(Include Area Code 152-3286	) 22c. OFFICE 00	SYMBOL	
2021	220 11:22 02		- 1 111		CCCURITY	CI ACCIDICATIO	N OF THIS PAGE	

i

SECURITY CLASSIFICATION OF THIS PAGE		
		1
		ĺ
		ļ
		ł
}		
		Ì
	,	
	•	
		ş
		İ
		ļ
		İ
		ć
		Ì

# IDENTIFYING THE CIRCADIAN CYCLE IN HUMAN INFORMATION PROCESSING DATA USING PERIODICITY ANALYSIS: A SYNOPSIS

#### ABSTRACT

Data from human information processing tasks frequently cannot satisfy the assumptions of many common periodicity techniques. This memorandum identifies appropriate techniques for data that cannot meet all of the assumptions of the more familiar ones.

### Acknowledgments

I would like to thank Dr. Paul Naitoh for his substantive comments on earlier versions of this memorandum and LCDR Dennis Reeves for his help in completing this project. I would also like to thank Kathleen Mayer for her valuable contributions in editing this manuscript.

Accesion For	4
NTIS CRA&I LTIC TAB Uninnounced Justification	
Py De You de la Francisco	
the state of	ric des
A-1	: / or 



#### INTRODUCTION

Historically, most studies examining the magnitude of the circadian effect on human information processing analyzed their data either by interpreting graphs or conducting analyses of variance (ANOVAs). Today, a disproportionate number of these studies still analyze their data using these methods rather than using some type of periodicity analysis (techniques that identify specific cycles in the data), which would be more informative. This puzzling state of affairs occurs because most studies of human information processing cannot meet the assumptions of many periodicity techniques. This memorandum describes some of the available techniques and the assumptions that must be met to use each technique. Problems in satisfying these assumptions are also discussed for repeated-measures assessments of information processing tasks.

#### PERIODICITY ANALYSES

7.50

Several different mathematical techniques can identify periodicities in data. The techniques described below are appropriate when the investigator is only interested in certain specific periodicities, such as the circadian cycle. In other words, the investigator is not asking "What periodicities are present?" but rather "Do the data show evidence of an x-hour cycle?" where x can be any value.

The selection of an analytical technique depends in part on satisfying its assumptions. These assumptions may be grouped into three major categories: 1) the regularity of the data sampling, 2) the behavior of the component periodicities (stationary vs. nonstationary), and 3) the independence of the observations. The strictest combination of assumptions is that the data are equally spaced, the component periodicities are stationary, and the observations are independent. Typically, data from information processing tasks cannot satisfy one or more of these assumptions and, consequently, must be analyzed using other techniques, such as ANOVA. The remainder of this memorandum discusses analytical techniques as a function of the assumptions that must be satisfied.

# EQUISPACED DATA, STATIONARY COMPONENT PERIODICITIES, AND INDEPENDENCE OF OBSERVATIONS

A variety of mathematical techniques--including Fourier analysis, autocorrelation, power spectrum analysis, and periodic (trigonometric) regression analysis (1,2)--are available to analyze equispaced data that have stationary periodicities and independent errors of observation. In addition to identifying the periodicitles present in the data, Fourier analysis and periodic regression analysis provide phase information that allows the time of peak performance to be determined. These techniques also allow the investigator to calculate both the change in performance over a fixed period of time and the best-fitting equation (see 2 for several excellent examples). Common statistical tests can be performed on the results of all of these analyses. For example, the results of an autocorrelation analysis can be tested to determine if the putative oscillations observed at a given frequency are statistically significant (1).

The major drawback of these techniques is that the data must be obtained at equally spaced intervals. Presently, many physiological variables

can be monitored continuously. Information processing data cannot be obtained, however, at equally spaced intervals without depriving the subject of sleep or waking the subject during the sleep cycle. Because the sampling frequency is determined primarily by the period of the cycle to be detected (see the Other Periodicity Analysis Problems Section), investigators studying the circadian cycle usually must either wake the subject for testing or preven, the subject from sleeping.

Some investigators (i.e., 3) claim that subjects reach baseline levels of performance on information processing tasks a few minutes after being awakened. This claim is not universally accepted. Few investigators question the effect of sleep deprivation on performance although short periods (a few hours) of deprivation may not have a statistically significant effect on performance. From a strictly practical point of view, either method may be used to obtain data at equally spaced intervals if the investigator has some assurance that the method will not significantly distort the data. If the subject needs an impractical amount of time after awakening to regain baseline performance levels or if short periods of sleep deprivation affect performance, the data may be collected at unequal intervals using the appropriate techniques.

# UNEQUALLY SPACED DATA, STATIONARY PERIODICITIES, AND INDEPENDENCE OF OBSERVATIONS

Trigonometric regression is the most well known technique that can be used to analyze data collected at unequally spaced intervals. The data are assumed to be independent and to reflect stationary cycles. Fourier analysis can also be performed for data collected at unequally spaced intervals (2) although it quickly becomes cumbersome and doen not appear to be used often in chronobiology. Easy-to-follow examples of urigonometric regression analysis for data collected at unequal, as well as equal, intervals are available (4).

### EQUALLY SPACED DATA, NORSTATIONARY PERIODICITIES, AND INDEPENDENCE OF OBSERVATIONS

Arguably, the most serious assumption concerning the assessment of the circadian cycle in human information processing data concerns the period of the cycle; the period is assumed to be stationary during the experiment. Usually, the investigator can safely assume that the component periodicities to not vary during the course of the experiment, that is, the frequency of each of the component periodicities is constant. This assumption has been challenged recently (5) for investigations of sustained operations, cumulative sleep deprivation, jet lag, and shift work. Frazier et al. (6) provide an excellent example of a large-magnitude change in periodicity in a highly controlled study of sustained operations. If the assumption of stationary periodicities is unwarranted, complex demodulation is the only time-series technique commonly available that can analyze nonstationary periodicities. This technique is restricted to "semistationary time series with slowly changing phase or amplitude or both" (5, p. 627).

Basically, complex demodulation first detrends the data, removing changes in the mean performance that occur over time (5). Then, it analyzes the rhythmic components of the data. Estimates of the instantaneous power and instant; neous phase are available at each sampling point for each

frequency examined. (This is in contrast to Fourier analysis, which provides one estimate of phase and one estimate of power for the entire time series.) By comparing the instantaneous power and phase across sampling points, the investigator can determine the extent to which the component periodicities changed during the experiment. Complex demodulation also provides estimates of the percentage of variance accounted for by each frequency being examined. Thus, the investigator can determine what percentage of the variance in the data was accounted for by trends in mean performance, the circadian (24-h) cycle, a 12-h cycle, an 8-h cycle, et cetera. Complex demodulation is relatively new and is still being developed. It should, therefore, be viewed with some caution.

### UNEQUALLY SPACED DATA, MONSTATIONARY PERIODICITIES, AND INDEPENDENCE OF OBSERVATIONS

Currently, this situation cannot be analyzed using any commonly available technique. Sing et al. (7) have attempted to develop a version of complex demodulation that can use unequally spaced data. To date, this version has not been widely accepted because of its complexity. Nevertheless, advances in complex demodulation may allow the analysis of unequally spaced data with nonstationary periodicities at some time in the future.

### NONINDEPENDENCE OF OBSERVATIONS

Regardless of the spacing of the data points, all the analysis techniques mentioned above assume that the observations are independent. This assumption is clearly untenable in the majority of situations where each subject in an experiment is measured repeatedly on an information processing task; in these situations, intertrial correlations usually exceed 0.50 and may exceed 0.90. The problem mathematically then becomes how to take these intercorrelations into account.

Unfortunately, analyzing such data, especially when they are collected at unequal intervals, is difficult. Nonstationary periodicities only worsen the problem. Currently, no analysis technique appears to be available for such data. Instead, such data are often analyzed by calculating the best fitting equation to one subset of the data and then crossvalidating the fit on the second subset (P. Naitoh, Naval Health Research Center, San Diego, CA, personal communication, September 8, 1988). Although primitive, this technique may provide sufficient information about the presence of periodicities for the investigator's purposes.

### OTHER PERIODICITY ANALYSIS PROBLEMS

Two other data analysis problems should be discussed. The first concerns the amount of data to be collected. The well-known Nyquist sampling theorem states that a signal must be sampled at a frequency of 2x

I Much repeated-measures human performance date are analyzed using analysis of variance (ANOVA) rather than a time series technique. In such experiments, the assumption of independence of errors of observation is also not tenable. To compensate for the violation of this assumption, criteria concerning the homogeneity of the variance-covariance matrix must be met.

to detect a periodicity with a frequency of x (8). That is, an investigator must sample twice as fast as the fastest frequency to be detected. Thus, to detect a circadian effect, an investigator theoretically must sample the variables of interest only twice during a 24-h period. Most investigators, however, assess performance at least four times during a 24-h period because of problems, such as the shifting cycle length, and the variance of human data.

A second related problem concerns the number of cycles (replications) to be sampled. For economic reasons, an investigator collects the minimum amount of data necessary to detect the periodicities in the dependent variable. At least five cycles may be necessary to permit a good estimate of an underlying periodicity (9). Estimates of a periodicity can be obtained using fewer than five cycles, but such estimates may be relatively inaccurate.

#### DISCUSSION

As noted in the introduction, many investigators apparently have been reluctant to use periodicity analysis on human information processing data. One possible explanation for this reluctance is that the assumptions and the requirements underlying some of the analyses cannot be met. The need for equally spaced intervals may be the most easily satisfied requirement. The assumption concerning independence of observations may be the most difficult to meet unless the investigator forgoes a repeated-measures design. The assumption concerning cycles of constant periods has only recently been challenged, and investigators have not yet identified the situations in which the "circadian" cycle shortens or lengthens significantly. Thus, for the immediate future, many investigators will continue to rely on graphs and ANOVAs to detect the presence of the circadian cycle in information processing data. As new periodicity techniques with less restrictive assumptions become available, investigators may shift the emphasis of their work from detecting differences in performance at various testing times to a more detailed examination of periodicities in the data.

#### REFERENCES

- 1. Oatley, K. and Goodwin, B., "The Explanation and Investigation of Biological Rhythms." In W. Colquhoun (Ed.), Biological Rhythms and Human Performance, Academic Press, London, UK, 1971, pp. 1-38.
- 2. Bliss, C., Statistics in Biology, Vol. 2, McGraw Hill Publishing Co., New York, NY, pp. 219-287, 1970.
- 3. Klein, K., Wegmann, H., and Hunt, B., "Desynchronization of Body Temperature and Performance Circadian Rhythm as a Result of Outgoing and Homegoing Transmeridian Flight." Aerospace Medicine, Vol. 43, pp. 119-132, 1972.
- 4. Naitoh, P., Englund, C., and Ryman, D., Circadian Rhythms Determined by Cosine Curve Fitting: Analysis of Continuous Work and Sleep Loss Data, Report No. 85-31, Naval Health Research Center, San Diego, CA, July 1985.

- 5. Sing, H., Thorne, D., Hegge, F., and Babkoff, H., "Trend and Rhythm Analysis of Time-series Data using Complex Demodulation." Behavior Research Methods, Instruments, and Computers, Vol. 17, pp. 623-629, 1985.
- 6. Frazier, T., Rummel, J., and Lipscomb, H., "Circadian Variability in Vigilance Performance." Aerospace Medicine, Vol. 39, pp. 383-395, 1968.
- 7. Sing, H., Hegge, F., and Redmond, D., "Complex Demodulation--Technique and Application." In E. Haus and H. Kabet (Eds.), Chronobiology 1982-1983, Karger, Basel, Switzerland, 1984, pp. 63-72.
- 8. Shannon, C., "A Mathematical Theory of Communication." Bell System Technical Journal, Vol. 27, pp. 379-423, 1948.
- 9. Deprins, K., Cornelisoon, G., and Malbecq, W., "Statistical Procedures in Chronobiology and Chronopharmacology." In A. Renberg, M. Smolensky, and G. Labrecque (Eds.), Annual Review of Chronopharmacology, Vol. 2, Pergamon, Oxford, UK, 1986, pp. 27-141.

### Other Related NAMEL Publications

- Damos, D.L., Some Considerations in the Design of a Computerized Human Information Processing Eattery, NAMRL Monograph 35, Naval Aerospace Medical Research Laboratory, Pensacola, FL, December 1987.

  (AD A199 491)
- Damos, D., A Review of Circadian Effects on Selected Human Information Processing Tasks, NAMRL Monograph 37, Naval Medical Research Laboratory, Pensacola, FL, April 1989.

<sup>\*</sup> These publications are available from DTIC, Cameron Station, Alexandria, VA 22314 (Phone: (C) 202/274-7633 or (AV) 284-7633).